

Expression of the 2'-N-acetyltransferase gene in *Providencia stuartii* is controlled by multiple *trans* acting loci and cell to cell communication.

Philip N. Rather, David Macinga and Elizabeth Orosz. Departments of Medicine and Molecular Biology and Microbiology, Case Western Reserve University and Veterans Affairs Medical Center, Cleveland, Ohio.

The chromosomal 2'-N-acetyltransferase gene [*aac(2')-Ia*] is universally present in *P. stuartii* and encodes resistance to aminoglycosides. In wild-type *P. stuartii*, *aac(2')-Ia* transcription occurs at low levels. We have utilized transposon mutagenesis to identify insertional mutations resulting in constitutive, high-level *aac(2')-Ia* expression. These insertions have identified three *trans* acting loci, designated *aarA*, *aarC*, and *aarD*, which either directly or indirectly regulate *aac(2')-Ia* expression. In addition, *aac(2')-Ia* expression is also regulated by an extracellular factor, termed AR-factor, which acts in a cell to cell signaling pathway to decrease *aac(2')-Ia* expression. AR-factor is a small (below 3 Kd), heat and pronase resistant molecule, which begins to accumulate in cell supernatants at mid-log phase. Growth of *P. stuartii* in the presence of high AR-factor concentrations resulted in an 8-fold decrease in *aac(2')-Ia* expression. Although mutations which prevent the synthesis of AR-factor would be predicted to have increased *aac(2')-Ia* expression, the *aarA*, *aarC* and *aarD* mutants all produce wild-type levels of AR-factor. Interestingly, *aarD* mutants are unable to grow in the presence of AR-factor, suggesting this extracellular factor plays a critical role in *P. stuartii* growth under certain conditions.